

The independent evaluation of soybean, dry bean, field pea and faba bean varieties found within this publication were made possible by your continued support through the Manitoba Pulse & Soybean Growers (MPSG) check-off. The objective of these trials is to provide the Manitoba pulse and soybean industry with independent, scientific information on variety performance and agronomic characteristics.

Field pea variety evaluations were coordinated with the Saskatchewan Regional Variety Testing Program. Field pea and faba bean variety evaluations were conducted by MCVET and partially sponsored by MPSG.

SOYBEANS

Herbicide tolerant soybean varieties were evaluated at 12 locations in 2020, reported by eastern and western regions in Manitoba. In eastern Manitoba, there are short-, mid- and long-season locations.

Mid- to long-season sites included Carman, Morris, Portage la Prairie and St. Adolphe. These sites are also referred to as core sites due to testing of all varieties at these locations. Shortseason sites included Arborg, Beausejour and Stonewall, where early- and midseason varieties were tested. In western Manitoba, sites included Boissevain, Dauphin, Hamiota, Melita and Swan River.

Herbicide tolerant first-year entry trials were also carried out at six of the 12 sites, including Boissevain, Carman, Hamiota, Melita, Morris and St. Adolphe.

This publication features the results from MPSG-sponsored trials.

Contents of this publication can only be reproduced with the permission of MPSG.

Conventional (non-GM) soybean varieties were tested at all sites listed for eastern Manitoba and at Melita and Swan River.

All soybean varieties are reported by very early-, early-, mid- and long-season maturity zones. Western Manitoba trials do not host long-season varieties, as they are generally ill-suited to the region.

DRY BEANS

Variety evaluations were conducted under wide- (>24 inches) and narrowrow (<12 inches) trials, and are reported separately in this guide.

Wide-row trials were conducted at five locations — Carman, Morden, Melita, Portage la Prairie and Winkler.

Narrow-row trials were conducted at four locations – Melita, Minto, Morden and Portage la Prairie.

Dry bean varieties are also reported by market class. These include navy, black, pinto, pink, Great Northern, dark red kidney, light red kidney and cranberry.

LENTILS

No lentil trials were conducted in Manitoba due to a lack of seed supply in the spring of 2020.

FIELD PEAS

Trials were conducted at eight locations in Manitoba, including Arborg, Boissevain, Carberry, Hamiota, Melita, Morden, Roblin and Swan River. Field pea varieties are reported by yellow, green, maple and forage market classes.

FABA BEANS

Registration trials were conducted at Roblin and Stonewall. Registered varieties

from these trials are reported by tannin and zero-tannin types.

USING THIS GUIDE

There are two types of data tables found in this guide – *Variety Descriptions* and *Yields by Location*. Variety description tables summarize long-term data, including maturity, yield and agronomic characteristics (e.g., disease resistance, lodging score). Yields by location tables summarize yield data from the current year at each location.

All variety trials were randomized with three replicates to allow for statistical analysis.

Statistical yield differences can be evaluated using only single-site year data, found in all *Yields by Location* tables. To compare yields, look at the least significant difference (LSD) value at the bottom of these tables. The LSD value represents the yield quantity (%) by which two varieties must differ, to conclude with 95% confidence that a true yield difference exists due to genetics.

For more information on how to use these tables, refer to the variety table keys in each section.

We acknowledge the contributions of all companies that submitted varieties and partners involved in planting, maintenance, note-taking, harvesting and data organization. Special thanks to staff at Manitoba Agriculture and Resource Development, AAFC, WADO, PCDF, PESAI, CMCDC and the private research companies that play an integral role in making this publication possible.

Key for All Variety Tables

Yield % Check – The average yield across all site years that the variety has been tested, relative to the check variety.

Site Years Tested – The total number of individual site years that a variety has been tested. For example, if a variety was tested at five sites for two years, the total site years would be 10. The greater the number, the more a variety has been tested under a greater range of environments. A variety is typically tested at two to five sites per year.


TSW (g/1000 seeds) – The thousand seed weight, referring to the seed weight in grams per 1000 seeds.


Resistance Rating – VG = very good G = good F = fair P = poor VP = very poor

CV % – The coefficient of variation (CV) is the statistical measure of random variation in a research trial. A CV of less than 15% generally indicates a more uniform trial and conclusive data.

LSD % – The least significant difference (LSD) is the quantity by which two varieties must differ to conclude with 95% confidence that a true difference exists due to genetics.

Sign. Diff. – The indication of whether significant differences were found between varieties. Yes = at least one variety is significantly different from another within one site. No = varieties are not significantly different within one site.







Maturity Zone	CHU	FFP (days)	Maturity Group
V. Early	<2250	<110	<00.2
Early	2250-2400	110–118	00.2-00.3
Mid	2401-2550	119–125	00.4-00.6
Long	>2550	>125	>00.6
I	-		

This map is based on 1981–2010 Climate Normal Data for cumulative Corn Heat Units (CHU, May 15 – Sept 20) and average frost-free period (FFP, days Tmin > 0°C). The map outlines the longest maturity suggested

The map outlines the longest maturity suggested for each production area, but earlier varieties can also perform well. Use in conjunction with the *Soybean Variety Guide*, which outlines varieties according to maturity zones. Manitoba Maturity Zone - Soybean varieties are organized into four maturity zones - very early-, early-, mid- and long-season. These categories reflect the Manitoba Soybean Maturity Zones map (page 2), based on long-term heat unit and frost-free period data. Varieties fit into respective zones based on average relative days to maturity. Each zone indicates the longest season varieties that should be selected for a given region.

Company Maturity Group - The maturity ranking provided by seed suppliers, indicating growing season length. Triple zero (000) and double zero (00) soybean varieties are best suited to Manitoba. Varieties currently tested in Manitoba range from 000 (earliest) to 0.1 (longest).

Type

E3 = Enlist E3[®] soybeans with 2,4-D choline, glyphosate and glufosinate herbicide tolerance.

RR1 = Roundup Ready 1 soybeans with glyphosate herbicide tolerance. R2Y = Genuity[®] Roundup Ready 2 Yield[®] soybeans with glyphosate herbicide tolerance.

R2X = Roundup Ready 2 Xtend® soybeans with dicamba and glyphosate herbicide tolerance.

DTM +/- Check - The number of days from planting to full maturity (R8 or 95% brown pod). It is expressed as + or - days relative to the check variety. Actual days to maturity (DTM) for the check variety is found in the shaded area at the bottom of the table. Average DTM is calculated from multiple site years. It is important to use long-term data for variety selection, as maturity can vary by year.

Hilum Colour - The hilum is the area of a soybean seed that was previously attached to the pod. Hilum colour is a marketing factor that varies among soybean varieties. Hilum colour can be clear (CL), yellow (Y), imperfect yellow (IY), grey (GR), light brown (LB), brown (BR), tan (TN), imperfect black (IB) or black (BL).

IDC Rating and Group - The iron deficiency chlorosis (IDC) rating is the severity of IDC expressed in a given variety on a 1-5 scale. The IDC group indicates the overall level of tolerance. Each year, ratings are conducted during the V2 to V3 stages at a site near Winnipeg that is prone to IDC. If a field is at moderate to high risk of IDC (Table 1), select a variety with a low (tolerant) rating.

IDC Ratings

1 = green leaves 2 = yellowish leaves 3 = green veins with yellow leaves

4 = brown dead tissue between green veins 5 = severe chlorosis and a stunted growing point soil test levels.

Soluble Salt		Carbonate (%)	
(mmhos/cm)	0 to 2.5	2.6 to 5	>5.0
0 to 0.25	Low	Low	Moderate
0.26 to 0.50	Low	Moderate	High
0.50 to 1.0	Moderate	High	Very high
>1.0	High	Very high	Extreme

Source: Agvise Laboratories

IDC Groups

T = tolerant ST = semi-tolerant S = susceptible

SCN - Variety resistance to soybean cyst nematode (SCN). The presence of SCN was confirmed for the first time in Manitoba in 2019. For full details of SCN findings, visit manitobapulse.ca.

PRR – Phytophthora root rot (PRR) race-specific resistance genes for each variety. Resistance genes that correspond with prevalent races in Manitoba are listed in Table 2. A new pathotype was most prevalent in Manitoba in 2018, according to Agriculture and Agri-Food Canada research. Soybean varieties with the rps 6 gene are resistant to this new pathotype.

Table 2. Resistance to *Phytopthora sojae* (rps) genes currently available in Manitoba for control of Phytophthora root rot.

Race of			Rps Gene		
P. sojae	1a	1c	1k	3a	6
New Pathotype	S	S	S	S	R
25	S	S	S	R	R
4	S	S	R	R	R
28	S	R	S	R	R
3	S	R	R	R	R

S = susceptible R = resistant

Source: Debra McLaren, AAFC



IDC Rating 1





IDC Rating 2.5

IDC Rating 3.5



IDC Rating 4.0

HERBICIDE TOLERANT SOYBEANS • VARIETY DESCRIPTIONS • EASTERN MANITOBA

		ICIDE IULEKANI									
Manitoba	Company							IC	00	Resis	tance
Maturity Zone	Maturity Group	Variety	Туре	DTM +/- Check†	Yield % Check	Site Years Tested	Hilum Colour	Rating (1—5)	Group	SCN	PRR
	000.7	Fresco R2X	R2X	-10	79	6	BL	2.2	ST	-	_
ery Early-	00.1	B0011RX	R2X	-9	83	6	TN	1.7	Т	-	1k
Season	000.5	NSC Wynyard RR2X	R2X	-9	78	6	BL	2.3	S	-	1a
Zone	00.1	S001-D8X	R2X	-6	95	6	IY	2.0	ST	-	1c
	00.2	Devo R2X	R2X	-6	90	17	BR	1.9	ST	-	-
	000.9	SI 000919XT	R2X	-5	89	6	BL	1.7	Т	-	-
	00.1	Torro R2	R2Y	-5	96	15	BL	2.2	ST	-	-
	000.9	RX000918	R2X	-5	97	11	BL	1.8	ST	yes	1c
	00.1	P001A48X	R2X	-4	96	11	TN	1.7	Т	-	1c
	00.4	NSC Culross RR2X	R2X	-4	100	11	BL	1.7	Т	-	1c
	00.2	NSC Redvers RR2X	R2X	-4	94	11	BL	1.9	ST	yes	1c
Early-	00.1	SI 001XTN	R2X	-3	102	17	BL	1.7	Т	yes	1k
Season	00.1	Prince R2X	R2X	-3	92	17	BL	1.8	ST	-	1k
Zone	00.3	Akras R2	R2Y	-3	102	21	BL	1.7	Т	-	1c
	00.4	TH89004 R2X	R2X	-3	89	11	BR	1.8	ST	-	1c
	00.5	S007-Y4	R2Y	-2	105	21	IY	2.0	ST	-	1c
	00.6	S006-M4X	R2X	-2	98	11	IY	1.9	ST	-	1c
	00.3	S003-Z4X	R2X	-2	102	6	BF	1.9	ST	-	1c
	00.3	Renuka R2X*	R2X	-2	102	6	LB	1.7	Т	yes	1c
	00.5	P005A83X	R2X	-1	100	11	BL	1.8	ST	yes	1c
	00.5	S005-C9X	R2X	-1	102	6	BL	2.3	S	-	1c
	00.6	RX Acron	R2X	-1	107	8	BL	1.8	ST	yes	-
	00.3	B0030L1	R2Y	-1	95	11	BR	1.9	ST	-	-
	00.3	Sunna R2X	R2X	0	103	17	GR	1.7	Т	yes	1c
	00.4	Bourke R2X	R2X	0	103	17	BL	1.8	ST	-	1k
	00.4	Merritt R2X	R2X	0	104	6	BI	1.9	ST	yes	1c, 1
	00.5	Foote R2	R2X	0	96	21	IY	1.8	ST	-	1c
	00.5	P005A27X	R2X	0	101	17	BR	1.9	ST	-	1c
	00.7	P007A90R*	RR1	0	100	22	BL	1.7	Т	yes	1c
	00.6	PS 0068 XR	R2X	0	104	12	BL	1.8	ST	-	1c
	00.6	NSC Sperling RR2Y	R2Y	0	107	17	IY	1.7	Т	-	1a, 3
	00.6	P006A37X	R2X	0	109	17	BR	1.8	ST	-	1c
Mid-	00.3	DKB003-29	R2X	0	98	21	BL	1.7	Т	yes	-
Season Zone	00.2	DKB002-32	R2X	0	103	6	BR	1.9	ST	yes	1k
Zone	00.6	NSC Cartier RR2X	R2X	1	105	6	BL	2.0	ST	-	3a
	00.4	B0040L1	R2Y	1	96	17	BR	1.7	Т	-	-
	00.5	Barker R2X	R2X	1	103	17	BL	1.8	ST	yes	1k
	00.3	TH 87003 R2X	R2X	1	96	21	BL	1.8	ST	yes	1c
	00.4	PV 16s004 R2X	R2X	1	101	17	BL	1.9	ST	yes	1k
	00.6	PV 19s006R2X	R2X	1	92	10	IB	2.0	ST	yes	1c
	00.5	DKB005-52	R2X	2	103	22	BL	1.8	ST	yes	1c
	00.7	S007-A2XS	R2X	2	110	6	GR	1.9	ST	-	-
	00.6	Kudo R2X	R2X	2	105	4	BL	1.7	Т	-	-
	00.7	P007A08X	R2X	2	110	5	GR	1.8	ST	-	1c
	00.6	B0066L1	R2Y	2	98	9	Y	1.9	ST	yes	1k
	00.7	Elmo E3	E3	2	107	4	BR	1.8	ST	yes	-
	00.7	RX00797	R2X	2	97	20	BL	1.7	Т	yes	1c
	00.7	PS 0074 R2	R2Y	3	107	17	BR	1.7	Т	-	-
	00.7	TH 88007R2X	R2X	3	102	18	BL	1.9	ST	-	1c
	00.7	PV 12s007 R2X	R2X	3	101	21	BL	1.9	ST	-	-
	00.5	TH 88005R2XN	R2X	3	100	18	BL	1.8	ST	yes	1c
	00.9	P00A49X	R2X	4	105	12	BR	1.7	Т	yes	1c
Long-	00.7	SI 007XTN	R2X	4	106	14	BL	1.8	ST	yes	1c
Season	00.6	DKB006-29	R2X	5	104	18	BL	1.7	Т	-	1k
Zone	00.8	NSC Winkler RR2X	R2X	5	106	12	BL	1.8	ST	yes	1c
	00.5	PV 10s005 RR2	R2Y	5	107	15	BL	1.9	ST	-	-
	00.9	NSC Aubigny RR2X	R2X	6	98	4	BL	1.6	Т	yes	1k
	00.9	TH89009 R2XN	R2X	6	108	8	BL	1.6	Т	yes	1k
	00.8	Astro R2	R2Y	7	110	8	BL	1.7	Т	-	1k
	00.8	Vidar R2X	R2X	8	104	14	BL	1.7	Т	yes	1c
	ACTEDICTICS										
IECK CHAF	ACTERISTICS	P007A90R		114	43	22					

† Maturity ratings were averaged across the Carman, Morris, Portage and St. Adolphe core sites over multiple years.

* 🖗 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

HERBICIDE TOLERANT SOYBEANS • YIELDS BY LOCATION • EASTERN MANITOBA

					2020 Yie	ld % Check		
Manitoba		DTM -	Early	Sites		Core	Sites	
Maturity Zone	Variety	DTM – +/- Check [†]	Beausejour	Stonewall	Carman	Morris	Portage	St. Adolphe
	Fresco R2X	-10	78	83	94	74	71	75
ery Early-	B0011RX	-9	63	97	109	67	74	102
Season	NSC Wynyard RR2X	-9	74	80	91	72	79	73
Zone	S001-D8X	-6	91	90	104	88	101	97
	Devo R2X	-6	86	87	96	77	89	90
	SI 000919XT	-5	64	83	116	85	98	90
	RX000918 P001A48X	-5 -4	80 91	102 99	121 107	89 92	99 88	92 103
	NSC Culross RR2X	-4	91	99	107	102	104	103
	NSC Redvers RR2X	-4	87	97	114	102	92	95
Early-	SI 001XTN	-3	83	108	123	99	100	111
eason Zone	Prince R2X	-3	80	93	111	92	85	97
Zone	Akras R2	-3	89	94	99	72	96	110
	TH89004 R2X	-3	72	92	108	89	91	82
	S007-Y4	-2	87	115	122	106	109	122
	S003-Z4X	-2	79	106	122	102	107	102
	Renuka R2X*	-2	91	102	118	98	108	96
	P005A83X	-1	79	97	115	104	98	113
	S005-C9X	-1	84	111	125	99	99	96
	RX Acron	-1	-	-	110	103	104	112
	B0030L1	-1	80	95	118	95	95	99
	Sunna R2X	0	84	107	113	104	112	103
	Bourke R2X	0	82	103	112	101	99	115
	Merritt R2X	0	98	108	109	98	102	118
	Foote R2	0	83	103	107	106	99	101
	P005A27X	0	106	88	100	95	99	107
	P007A90R	0	100	100	100	100	100	100
	PS 0068 XR	0	-	-	112	111	87	93
	NSC Sperling RR2Y	0	105	115	117	101	106	106
Mid-	P006A37X	0	109	122	123	113	117	112
eason Zone	DKB003-29	0	97	95	115	94	94	103
20110	DKB002-32	0	93	104	106	107	99	110
	NSC Cartier RR2X	1	99	105	123	107	93	103
	B0040L1	1	88	97	124 122	103 109	107 96	111
	Barker R2X TH87003 R2X		86	90	98	88	87	99
	PV 16s004 R2X	1	86	105	109	105	104	111
	PV 19s006R2X	1	93	101	100	89	98	97
	DKB005-52	2	100	104	129	107	105	113
	S007-A2XS	2	103	112	119	97	117	120
	Kudo R2X	2	-	-	104	99	107	111
	B0066L1	2	-	-	110	101	94	102
	Elmo E3	2	-	-	107	102	114	106
	RX00797	2	78	95	108	91	93	99
	PS 0074 R2	3	-	-	106	114	111	113
	TH 88007R2X	3	87	104	112	102	97	102
	PV 12s007 R2X	3	92	98	112	103	93	104
_ong-	TH 88005R2XN	3	95	99	122	104	92	92
eason –	P00A49X	4	-	-	120	99	111	106
Zone	SI 007XTN	4	81	98	131	104	91	103
	DKB006-29	5	-	-	127	103	108	107
	NSC Winkler RR2X	5	-	-	114	102	93	113
	TH89009 R2XN Vidar R2X	6	-	-	105	95 100	105 105	109 97
		0		_	112	100	105	97
	P007A90R	114	55	43	48	58	54	32
		DTM				u/ac		
		CV %	9.1	6.6	6.9	5.5	7.4	7.8
		LSD %	13	11	12	9	12	13
		Sign. Diff. Seeding Date	yes May 19	yes May 23	yes May 26	yes May 22	yes Jun 3	yes May 20

† Maturity ratings were averaged across the core sites over multiple years.

* 🖗 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

HERBICIDE TOLERANT SOYBEANS • VARIETY DESCRIPTIONS & YIELDS BY LOCATION • WESTERN MANITOBA

Manitoba	Company				-	1(00	Resis	tance		202	0 Yield % Ch	eck	-
Maturity Zone	Maturitý Group	Variety	DTM +/- Check†	Yield % Check	Site Years Tested	Rating (1–5)	Group	SCN	PRR	Boissevain	Dauphin	Hamiota	Melita	Swan Rive
	000.7	B00071RX	-13	85	7	1.7	Т	-	1k	78	77	104	96	85
	000.5	Amirani R2*	-9	90	7	1.8	ST	-	1k	90	91	109	94	80
	00.2	DKB002-32	-7	104	5	1.9	ST	yes	1k	99	103	118	103	94
Very Early-	000.7	Fresco R2X	-7	92	7	2.2	ST	-	-	80	86	115	94	103
Season	000.8	NSC Watson RR2Y	-6	94	27	2.1	ST	-	6	100	91	126	102	105
Zone	00.1	B0011RX	-6	108	5	1.7	т	-	1k	112	88	134	96	108
	000.5	NSC Wynyard RR2X	-5	95	5	2.3	S	-	1a	94	83	118	85	96
	000.8	NocomaR2*	-5	85	17	2.0	ST	-	1c	67	78	81	81	74
	000.9	S0009-M2	-5	95	27	2.0	ST	_	6	104	94	122	96	104
	000.9	RX000918	-3	96	12	1.8	ST	yes	1c	111	94	127	105	78
	000.5	DKB0005-44	-3	91	12	1.9	ST	yes	1c	93	83	107	98	93
	000.9	Fisher R2X	-3	97	7	1.8	ST	yes	1k	106	97	126	103	72
	00.4	TH89004 R2X	-2	101	7	1.8	ST	_	1c	114	95	115	96	96
	00.1	S001-D8X	-2	108	5	2.0	ST	_	1c	106	97	116	101	124
	00.1	Torro R2	-1	95	17	2.2	ST	_	_	107	90	116	101	95
	000.9	PV 15s0009 R2X	-1	99	11	2.1	ST	yes	1c	104	97	123	97	109
	00.3	P003A97X	-1	98	7	1.9	ST	yes	1k	90	100	120	97	88
	00.3	S003-Z4X	-1	107	7	1.9	ST	-	1c	100	102	134	104	106
	00.5	P001A48X	-1	100	7	1.7	T	_	1c	96	98	124	93	108
	00.3	Mahony R2	-1	100	29	2.9	S	-	-	106	96	123	101	
Early- Season	00.5	SI 001XTN	-1	97	12	1.7	T	yes	- 1k	85	104	116	106	79
Zone	00.5	S007-Y4	-1	102	30	2.0	ST	yes	1c	107	112	136	100	-
	00.3	Prince R2X	-1	97	11	1.8	ST	_	1k	98	95	129	103	107
				89	7				1K	84			93	78
	00.2	Devo R2X	-1			1.8	ST T				80	112	105	78
	00.3	Renuka R2X*	0	106	6	1.7			1c	106	100	129		-
	00.5	TH 88005R2XN	0	95	9	1.8	ST	yes	1c	100	103	98	100	-
	00.5	P005A83X	0	106	7	1.8	ST	yes	1c	104	105	126	100	112
	00.2	NSC Redvers RR2X	0	97	10	1.9	ST	yes	1c	118	92	125	106	-
	00.3	Akras R2	0	100	31	1.7	T	-	1c	100	100	100	100	100
	000.9	SI 000919XT	0	100	5	1.7	T	-	-	91	89	122	104	89
	000.9	DKB0009-89	0	99	12	1.9	ST	yes	1c, 1k	113	101	117	106	98
	00.3	NSC Newton RR2X	0	86	15	2.0	ST	-	-	93	85	103	92	-
	00.3	Sunna R2X	1	102	10	1.7	Т	yes	1c	98	104	135	106	-
	00.5	S005-C9X	1	112	4	2.3	S	-	1c	112	98	130	106	-
	00.3	TH 87003 R2X	1	95	18	1.8	ST	yes	1c	84	86	96	99	74
	00.3	B0030L1	2	94	7	1.9	ST	-	-	88	93	111	95	83
Mid-	00.4	Bourke R2X	2	103	6	1.9	ST	-	1k	104	107	116	101	-
Season Zone	00.6	P006A37X	2	108	11	1.8	ST	-	1c	107	112	124	107	110
Zone	00.6	PS 0068 XR	3	110	4	1.8	ST	-	1c	114	94	120	108	-
	00.5	P005A27X	3	100	12	1.9	ST	-	1c	85	98	111	103	102
	00.4	Merritt R2X	3	106	4	1.9	ST	yes	1c, 1k	91	103	120	108	-
	00.4	PV 16s004 R2X	3	99	10	1.8	ST	yes	1k	109	100	116	95	-
	00.5	Kudo R2X	3	102	6	1.7	Т	-	-	110	99	116	97	-
HECK CHAR	RACTERISTICS	Akras P2	174	54	31					48	45	55	66	44
		Akras R2	124 DTM	54 bu/ac	3 I site years					40	40	bu/ac	00	44
									CV %	13	4.3	4.4	3.8	8.2
									LSD %	21	7	8	6	13
									gn. Diff.	yes May 28	yes May 15	yes May 16	yes May 19	yes May 25
								seeair	ng Date	May 28	May 15	May 16	May 19	May 25

† Maturity ratings were averaged across the western sites over multiple years.

* 🖗 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

HERBICIDE TOLERANT SOYBEANS • YIELDS BY LOCATION • EASTERN FIRST YEAR ENTRY LEVEL

		SOIDERINS II			
Manitoba				2020 Yield % Check	
Maturity		DTM			
Zone	Variety	+/- Check [†]	Carman	Morris	St. Adolphe
	Halo R2X	-7	96	72	86
Very Early-	Aveta R2X	-6	113	77	60
Season Zone	Experimental lines that are	being tested/proposed for	registration in Canada		
Lone	CFS20.1.R2	-9	86	58	70
	SI 000920XT	-5	117	86	92
Early-	Young R2X	-5	121	91	90
Season – Zone –	Hart R2X	-3	131	109	98
10.110	Mikado R2X	-2	106	99	80
	PV 22S002 R2X	-1	116	99	101
	P007A90R	0	100	100	100
Mid-	Mao R2X	2	122	108	101
Season	Experimental lines that are	being tested/proposed for	registration in Canada		
Zone	CFS20.2.R2	-1	106	80	98
	PV EXP 20-S4	0	130	103	107
	EXP00520XR	2	122	97	95
	SI 00820XTN	3	126	105	92
	NSC EXP004X	3	131	103	80
	SI 0220XT	4	110	91	109
Long- Season	TH81007 R2XN	4	121	99	88
Zone	Bronco R2X	4	102	86	100
	SI 0120XTN	5	119	108	85
	Experimental lines that are	being tested/proposed for	registration in Canada		
_	TH79009E	3	127	108	104
CHECK CHAR					
	P007A90R	115	43	59	35
		DTM		bu/ac	
		CV 9 LSD 9		5.3	6.5
				8	10
		Sign. Dif Seeding Date		yes May 22	yes May 20
		Harvest Date		Sep 30	Sep 22

† Maturity ratings were averaged across the Carman, Morris and St. Adolphe sites.

HERBICIDE TOLERANT SOYBEANS • YIELDS BY LOCATION • WESTERN FIRST YEAR ENTRY LEVEL

Manitaha			2020 Yield	% Check
Manitoba Maturity Zone	Variety	DTM +/- Check [†]	Hamiota	Melita
	DKB0003-24	-6	115	91
	Mynarski R2X	-6	118	97
Very Early-	Halo R2X	-5	119	90
Season Zone	Experimental lines that are	being tested/proposed for registration	in Canada	
20110	PV EXP 20-S2	-5	123	93
	PV EXP 20-S1	-4	113	95
	NSC Watson RR2Y	-2	111	96
	Inferno R2X	-2	106	62
	S0009-F2X	-2	122	100
	SI 000920XT	-2	116	88
Early-	Aveta R2X	-1	108	90
Season – Zone –	Major R2X	0	128	95
20110	Akras	0	100	100
	Experimental lines that are	being tested/proposed for registration	in Canada	
	CFS20.1.R2	-1	86	86
	CFS20.2.R2	-1	108	90
	Hart R2X	2	135	101
Mid-	DKB0008-87	3	128	104
Season Zone	Mikado R2X	3	118	103
20.10	PV 225002 R2X	3	103	95

HERBICIDE TOLERANT SSOYBEANS • YIELDS BY LOCATION • WESTERN FIRST YEAR ENTRY LEVEL continued

Manitoba			2020	Yield % Check
Maturity Zone	Variety	DTM +/- Check [†]	Hamiota	Melita
	Young R2X	3	114	98
	Bronco R2X	6	88	86
Mid-	Experimental lines that a	re being tested/proposed for registration in Canad	a	
Season Zone	CFS20.3 R2	3	111	100
Zone	PV EXP 20-S4	5	106	100
	EXP00520XR	5	109	96
CHECK CHAR	ACTERISTICS			
	Akras R2	108	52	64
		DTM		bu/ac
		CV %	5.8	4.4
		LSD %	11	7
		Sign. Diff.	yes	yes
		Seeding Date	May 16	May 19
		Harvest Date	Oct 1	Sep 18

† Maturity ratings were averaged across the Hamiota and Melita sites.

CONVENTIONAL SOYBEANS VARIETY DESCRIPTIONS

Manitoba	Company						IC	C
Maturity Zone	Maturity Group	Variety	DTM +/- Check†	Yield % Check	Site Years Tested	Hilum Colour	Rating (1–5)	Group
	000.8	Norfolk	-6	94	20	IY	2.3	S
Early- Season	000.7	Fjord	-4	95	12	IY	1.9	ST
Zone	000.9	AAC Halli*	-3	101	39	Y	2.1	ST
Zone	00.2	Siberia	-2	110	11	IY	1.9	ST
	00.3	OAC Prudence	0	100	136	Y	1.6	Т
	00.3	Maxus	0	99	18	IY	2.0	ST
	00.3	Reynolds	2	107	17	IY	2.3	S
	00.4	Liska*	4	118	6	IY	2.3	S
	00.6	Kebek	6	100	17	Y	1.8	ST
Mid-	00.7	Abaca	6	127	3	IY	1.8	ST
Season	00.9	DH863	6	96	25	IY	2.3	S
Zone	Experiment	al lines that are being test	ed/proposed for regis	tration in Canada				
	000	SVX21T000S1	1	110	6	IY	2.1	ST
	00	SVX21T00S2	3	108	6	IY	1.7	Т
	00.7	CER10-11.97	5	101	6	IY	1.9	ST
	00.8	CER14-640	5	106	6	IY	2.1	ST
	00.3	PR130167Z1-02	6	111	3	BR	1.8	ST
	00.8	Meteor	7	100	17	IY	2.3	S
	00.8	Aurelina	8	116	3	IY	1.9	ST
1	00.6	Maya*	9	101	3	IY	1.7	Т
Long- Season	00	Stanley	10	113	6	IY	2.1	ST
Zone	0.3	Astor	12	114	12	Y	2.0	ST
Zone	Experiment	al lines that are being test	ed/proposed for regis	tration in Canada				
	00.7	CER14-142	9	111	6	Y	2.1	ST
	00.9	DL18.3004	11	117	6	Y	2.1	ST
IECK CHAI	RACTERISTICS	OAC Prudence	112	47	136			
			DTM	bu/ac	site years			

† Maturity ratings were averaged across the core sites over multiple years.

Pulse Soybean GROWERS Working for You

MANITOBA

* 😡 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.



For more information visit manitobapulse.ca

Follow us 👎 У O 🕨 YouTube

CONVENTIONAL SOYBEANS • YIELDS BY LOCATION • EASTERN MANITOBA

		_			2020 Yield	l % Check		
Manitoba		_		Early Sites		Core Sites		
Maturity		DTM						
Zone	Variety	+/- Check [†]	Arborg	Beausejour	Stonewall	Carman	Morris	Portage
E e el e	Norfolk	-6	84	63	85	97	95	106
Early- Season	Fjord	-4	92	87	87	111	100	92
Zone -	AAC Halli*	-3	90	89	113	100	98	103
Zone	Siberia	-2	106	97	95	121	116	115
	OAC Prudence	0	100	100	100	100	100	100
	Maxus	0	-	-	-	106	103	98
	Reynolds	2	94	94	109	107	107	100
	Liska*	4	-	-	-	121	118	110
	Kebek	6	99	87	92	111	118	107
Mid-	Abaca	6	-	-	-	133	134	117
Season	DH863	6	-	-	-	99	99	87
Zone	Experimental lines	that are being tested/	proposed for reg	istration in Canada				
	SVX21T000S1	1	112	97	104	132	103	112
	SVX21T00S2	3	105	108	105	126	105	103
	CER10-11.97	5	104	103	99	99	108	91
	CER14-640	5	103	110	105	103	112	100
	PR130167Z1-02	6	-	-	-	124	110	103
	Meteor	7	100	102	99	96	97	88
	Aurelina	8	-	-	-	127	117	108
	Maya*	9	-	-	-	105	102	96
Long-	Stanley	10	-	-	-	109	112	91
Season	Astor	12	-	-	-	104	110	97
Zone	Experimental lines	that are being tested/	proposed for reg	istration in Canada				
	CER14-142	9	110	116	116	110	121	95
	DL18.3004	11	-	-	-	114	112	99
HECK CHAR	ACTERISTICS		281					
	OAC Prudence	112	44	52	32	40	51	52
		DTM			bu	/ac		
		CV %	7.1	9.8	6.6	8.8	4.4	6.9
		LSD %	12	16	-11	16	8	11
		Sign. Diff.	yes	yes	yes	yes	yes	yes
		Seeding Date	May 15	May 20	May 23	May 26	May 22	Jun 3
		Harvest Date	Sep 29	Oct 7	Sep 24	Oct 1	Sep 29	Oct 6

† Maturity ratings were averaged across the core sites over multiple years.

* 🖗 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

CONVENTIONAL SOYBEANS • YIELDS BY LOCATION • WESTERN MANITOBA

Manitoba						2020 Yie	ld % Check
Maturity Zone	Variety	DTM +/- Check†	Yield % Check	Site Years Tested	Hilum Colour	Melita	Swan River
Early-	Ambella	-9	98	2	BR	95	103
Season	AAC Halli*	-3	93	6	Y	103	82
Zone	Fjord	-3	95	6	IY	95	105
	Siberia	-2	103	4	IY	102	109
	Liska*	-1	111	2	IY	100	112
	OAC Prudence	0	100	9	Y	100	100
Mid-	Maxus	0	95	6	Y	93	100
Season Zone	Maya*	4	97	2	IY	88	89
Zone	Experimental lines th	nat are being tested/pr	oposed for registration	on in Canada			
	PR130933Z-05	-1	92	2	Y	95	89
	PR130167Z1-02	4	90	2	BR	98	81
HECK CHAR	ACTERISTICS						
	OAC Prudence	118	36	9		39	31
		DTM	bu/ac	site years		bu	u/ac
					CV %	4.1	9.2
					LSD %	7	15
					Sign. Diff.	yes	yes
					Seeding Date	May 21	Jun 4
					Harvest Date	Sep 17	Oct 13

† Maturity ratings were averaged across the Melita and Swan River sites over multiple years.

* 🔞 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

Key for Field Pea Variety Tables

Maturity Ratings – early = -3 days relative to the check mid = -2 to -1 days late = 0 to +3 days

 $\label{eq:relative_state} \begin{array}{ll} \mbox{Relative Vine Length} - S = short & M = medium & L = long \\ \mbox{VL} = very \mbox{ long} \end{array}$

Green Seed Coats – G = 0-10% green seed coats F = 11-25% green seed coats

Seed Coat Dimpling – VG = 0–5% of seeds dimpled G = 6-20% of seeds dimpled F = 21-50% of seeds dimpled

Bleaching – The resistance rating of green pea to bleaching. Bleaching does not apply to other market classes of peas, indicated by n/a.

Mycosphaerella Blight – All pea varieties listed have "fair" resistance to Mycosphaerella (Ascochyta) blight.

Fusarium Wilt – Varieties with good resistance to one strain of fusarium wilt may be susceptible to other strains.

									Resistance			-
Market Class/Variety	Maturity Rating†	Yield % Check	Site Years Tested	Relative Vine Length	TSW (g/1000 seeds)	Green Seed Coats	Seed Coat Breakage	Seed Coat Dimpling	Seed Coat Bleaching	Lodging	Powdery Mildew	Fusarium Wilt
YELLOW												
AAC Ardill	mid	99	35	М	240	G	G	G	n/a	G	VG	G
AAC Asher*	mid	102	12	S	260	G	F	F	n/a	G	VG	F
AAC Carver*	early	104	29	L	240	G	G	G	n/a	G	VG	F
AAC Chrome*	long	108	25	М	240	G	G	G	n/a	G	VG	F
AAC Delhi*	mid	102	13	М	290	G	F	F	n/a	G	VG	F
AAC Lacombe**	long	100	33	L	270	F	F	G	n/a	G	VG	F
AAC Profit*	long	101	12	М	230	G	F	G	n/a	G	VG	F
CDC Amarillo	long	100	35	М	230	G	F	F	n/a	VG	VG	G
CDC Athabasca*	long	96	19	L	300	G	F	F	n/a	VG	VG	G
CDC Canary*	early	98	19	L	230	F	G	F	n/a	VG	VG	F
CDC Inca*	mid	104	33	L	230	F	G	G	n/a	G	VG	F
CDC Lewochko*	mid	103	19		230	G	G	G	n/a	VG	VG	F
CDC Meadow	early	97	85	м	220	G	G	G	n/a	G	VG	F
CDC Saffron	mid	97	49	М	250	G	G	F	n/a	G	VG	F
CDC Spectrum*	long	95	19	L	240	G	G	G	n/a	VG	G	F
GREEN		_										
AAC Comfort*	long	97	24	м	260	n/a	G	G	G	G	VG	F
Blueman*	long	96	6	м	230	n/a	VG	G	F	G	VG	F
CDC Forest*	long	101	19	L	230	n/a	G	G	G	G	VG	F
CDC Greenwater	mid	97	34	м	220	n/a	VG	G	G	G	VG	G
CDC Limerick	long	96	34	М	210	n/a	VG	G	G	VG	VG	F
CDC Spruce*	long	98	19		240	n/a	F	F	G	G	VG	F
CDC Striker	early	87	89	м	230	n/a	VG	G	G	VG	Р	G
MAPLE	,											
AAC Liscard	early	93	24	М	180	n/a	_	_	n/a	G	VG	_
FORAGE	,											
DL Delicious	long	73	6	VL	220	_	VG	F	n/a	F	-	-
DL Goldeye*	long	69	13	VL	160	G	VG	VG	n/a	VP	-	-
DL Lacross	mid	86	13	VL	190	F	VG	G	n/a	G	-	-
CHECK CHARACTERIST												
CDC Amarillo	94	76	35									
	DTM	bu/ac	site years									

† Maturity ratings were averaged across Hamiota, Melita, Morden and Swan River. * 🕐 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991. ** 🛞 Indicates a variety that is protected by Plant Breeder's Rights legislation that complies with UPOV 1991.

FIELD PEAS YIELDS BY LOCATION

	2020 Yield % Check									
Market Class/Variety	Carberry	Hamiota	Melita	Morden	Roblin	Swan River				
YELLOW										
AAC Ardill	112	101	98	109	125	109				
AAC Asher	106	88	108	118	100	109				
AAC Carver*	107	91	100	110	116	102				
AAC Chrome*	116	85	107	121	113	114				
AAC Delhi*	95	94	105	117	111	110				
AAC Lacombe**	109	90	98	122	106	109				
AAC Profit*	112	93	98	100	111	109				
CDC Amarillo	100	100	100	100	100	100				

	2020 Yield % Check											
Market Class/Variety	Carberry	Hamiota	Melita	Morden	Roblin	Swan River						
CDC Athabasca*	110	94	97	90	101	103						
CDC Canary*	113	99	91	102	115	102						
CDC Inca*	118	106	101	112	104	111						
CDC Lewochko*	100	116	98	99	107	108						
CDC Meadow	100	97	100	108	88	106						
CDC Saffron	109	89	91	106	99	101						
CDC Spectrum*	89	90	94	98	95	104						
GREEN												
AAC Comfort*	122	89	101	97	115	107						
Blueman*	98	85	95	94	121	107						
CDC Forest*	112	102	101	86	102	107						
CDC Greenwater	107	98	93	101	95	105						
CDC Limerick	97	101	96	97	88	98						
CDC Spruce*	88	88	97	100	113	107						
CDC Striker	99	81	89	96	105	93						
MAPLE												
AAC Liscard	97	89	96	101	100	96						
FORAGE												
DL Delicious	95	66	67	89	78	70						
DL Goldeye*	82	71	57	61	85	66						
DL Lacross	98	86	95	103	93	71						
CHECK CHARACTERISTICS												
CDC Amarillo	94	67	94	78	72	92						
			bu	ı/ac								
CV %	10.2	4.5	7.2	6.8	11.0	4.8						
LSD %	17	7	11	11	19	8						
Sign. Diff.	yes	yes	yes	yes	yes	yes						
Seeding Date	May 6	May 7	May 6	May 19	May 7	May 14						
Harvest Date	Aug 24	Aug 25	Aug 17	Aug 24	Aug 27	Aug 27						

* 🚱 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

** 🛞 Indicates a variety that is protected by Plant Breeder's Rights legislation that complies with UPOV 1978.

Key for Faba Bean Variety Table

Tannin vs. Zero-Tannin Varieties – Tannin varieties with coloured flowers and tan-coloured seed coats cannot be fed directly to livestock. Zero-tannin varieties with white flowers and seed coats can be fed directly to livestock. **DTM** – The number of days from planting to swathing. Days to maturity (DTM) may vary depending on the planting date.

FABA BEANS • VARIETY DESCRIPTIONS AND YIELDS BY LOCATION

				_	2020 Yie	ld % Check
Market Class/ Variety	DTM +/- Check	Yield % Check	Site Years Tested	TSW (g/1000 seeds)	Roblin	Stonewall
COLOURED FLOWER (TANNIN)						
Fabelle*	0	100	1	533	100	100
CHECK CHARACTERISTICS						
Fabelle	105	3852	5		3768	4118
	DTM	lbs/ac	site years		lb	s/ac
WHITE FLOWER (ZERO TANNIN)						
Snowbird**	0	100	16	495	-	100
DL Rico*	5	89	3	566	-	93
DL Tesoro*	6	109	3	511	-	103
CHECK CHARACTERISTICS						
Snowbird	104	4824	16		-	3748
	DTM	lbs/ac	site years		lb	s/ac
				CV %	-	5.1
				LSD %	-	8
				Sign. Diff.	-	yes
				Seeding Date	May 7	May 17
				Harvest Date	Sep 10	Sep 8

* 🔞 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

** (Indicates a variety that is protected by Plant Breeder's Rights legislation that complies with UPOV 1978.

DTM +/- **Check** – The number of days from planting to full maturity (90% of plants ready for harvest). It is expressed as + or – days relative to the check variety. Actual days to maturity (DTM) for the check variety is found in the shaded area at the bottom of the table.

Lodging (1–5) – The lodging rating at harvest on a scale of one to five. The greater the value, the more lodged the crop. For example, 1 = standing upright, 5 = flat on the ground.

Plant Height (cm) – The distance measured from the soil surface to the top of the plant at flowering.

Pod Height (% >5 cm) – The visual estimation of the % of pods greater than 5 cm from the soil surface at harvest.

CBB Severity (0–5) – The average visual rating of common bacterial blight (CBB) on 10 plants per plot at the yellow pod (R7) stage.

- $\mathbf{0}=\mathbf{N}\mathbf{o}$ observable lesions or other signs of infection
- 1 = < 5% of plant area (leaf and stem hypocotyls) diseased
- 2 = 5-10% of plant area diseased
- 3 = 10-25% of plant area diseased
- 4 = 25-50% of plant area diseased

 $5=50{-}100\%$ of plant area diseased or death of seedling

CBB Incidence (%) – The average visual rating of % leaf tissue infected by CBB on 10 plants per plot at the R7 stage.

WM Incidence (%) – The average visual rating of the % of plants infected by white mould (WM) on 10 plants per plot at full maturity (R9).

DRY BEANS • VARIETY DESCRIPTIONS

		Yield	Site	TSW		Plant	Pod	CBB	CBB	WM
Market Class/Variety	DTM +/- Check	% Check	Years Tested	(g/1000 seeds)	Lodging (1–5)	Height (cm)	Height (% > 5 cm)	Severity (0–5)	Incidence (%)	Incidence (%)
NAVY	+/- T9905	% T9905		,	. ,	. ,				. ,
AAC Shock	-2	97	8	209	2	52	90	2	21	0
Bolt	-2	93	19	204	1	53	90	2	19	0
Indi	-2	101	28	172	1	55	92	3	19	0
AAC Argosy	-1	104	12	193	2	54	91	2	18	0
Armada	0	97	4	187	2	58	88	2	17	0
Nautica	0	89	18	163	2	50	93	2	20	0
T9905	0	100	36	199	2	52	90	2	15	0
HMS Medalist	1	99	6	192	2	57	88	3	17	0
SV1893GH*	in A	95	13	192	2	55	88	1	10	0
Varieties that are regis						55	00	I	10	0
S09-27C	-4	86	2	228	2	54	90	3	15	0
15094	-4	103	6	204		56	90 89	2	15	0
15094	3	103	6	204	3	58	88	3	25	0
		101	0	204	5	50	00	2	25	0
CHECK CHARACTERIST T9905	100	2366	36							
19903	DTM	lbs/ac	site years							
BLACK			site years							
CDC Blackstrap*	+/- Eclipse -5	% Eclipse 94	17	209	1	44	89	2	13	0
Ace	-2	99	6	195	2	55	92	3	17	0
CDC Jet	-2	89	43	195	1	50	92	2	17	0
CDC Superjet	-1	88	31	192	2	50	92	2	15	0
Black Tails	0	98	6	198	2	55	92	3	25	0
Eclipse**	0	100	45	197	1	54	90	3	20	0
Zenith	3	96	45	220	1	54	93	3	33	0
	-		-			21	93	3	33	0
Varieties that are regis W11-02-152	0	83	ed for registrat	234	a 2	58	88	3	25	0
		65	2	234	Z		00	2	25	0
CHECK CHARACTERIST	96	2449	45							
Eclipse	DTM	lbs/ac	45 site years							
PINK			site years							
Floyd	+/- Floyd O	% Floyd 100	27	340	4	46	63	3	42	0
CHECK CHARACTERIST		100	21	5-10	-7	-10	00	5	-†∠	0
Floyd	92	2400	27							
noya	DTM	lbs/ac	site years							
PINTO	+/- Windbreaker	% Windbreaker	Site years							
SV6139GR*	-2	104	24	332	2	54	87	3	20	0
Vibrant	-2	108	17	327	2	63	86	3	27	0
SV6533GR*	-1	97	9	423	3	55	78	3	27	0
Windbreaker	0	100	52	359	3	50	81	3	22	0
Cowboy*	0	108	6	375	2	63	88	3	22	0
	2	108	20	375	2	65	87	3	18	0
Monterrey	2	100	20	000	2	00	0/	S	10	0

DRY BEANS 🔸 VARIETY DI	ESCRIPTIONS continued									
Market Class/Variety	DTM +/- Check	Yield % Check	Site Years Tested	TSW (g/1000 seeds)	Lodging (1–5)	Plant Height (cm)	Pod Height (% > 5 cm)	CBB Severity (0—5)	CBB Incidence (%)	WM Incidenc (%)
La Paz	4	99	21	320	3	61	86	3	26	0
Varieties that are regis	tered in the US or be							-		-
16-NP1	4	86	4	323	3	64	88	2	20	0
19-382	5	89	4	431	3	53	86	2	20	0
ND Palomino*	5	106	6	387	4	59	82	3	23	0
19-279	7	96	4	396	3	66	87	2	17	0
CHECK CHARACTERIST				550	5	00	07	2	17	
Windbreaker	93	2657	56							
	DTM	lbs/ac	site years							
GREAT NORTHERN	+/- Pink Panther	% Pink Panther								
Aries	-2	135	13	370	3	50	87	3	30	0
Varieties that are regis	tered in the US or be	ing tested or propos	ed for registrat	ion in Canad	а					
Powderhorn	-3	133	8	363	3	47	84	3	25	3
DARK RED KIDNEY	+/- Pink Panther	% Pink Panther								
Red Hawk	4	65	15	525	2	36	86	3	28	0
Dynasty	6	96	6	509	3	61	84	3	26	0
Montcalm	6	81	5	460	2	47	86	4	32	0
Varieties that are regis	tered in the US or be	ing tested or propos								
161164	5	74	3	383	3	51	85	3	24	2
Red Rover	5	56	6	428	3	51	82	3	27	0
LIGHT RED KIDNEY	+/- Pink Panther	% Pink Panther		120			02			
Big Red	0	99	24	503	2	45	86	3	27	0
Pink Panther	0	100	56	513	2	50	88	3	30	0
Varieties that are regis	-					50	00	J	50	0
Red Dawn	-9	99	8	511	u 1	48	84	3	23	0
		,,,		511		U		5		
Pink Panther	99	1981	56							
	DTM	lbs/ac	site years							
CRANBERRY	+/- Etna	% Etna								
OAC Racer	-1	110	3	478	1	48	85	3	30	0
Etna	0	100	58	509	1	44	84	3	34	0
OAC Candycane	4	134	3	501	1	52	88	2	13	0
AAC Scotty	5	111	14	526	G	41	85	3	26	0
Varieties that are regis								5	20	÷
SV3709GC	-4	109	6	568	1	39	87	4	47	0
Amaranto	-3	101	3	459	1	47	87	3	30	0
CR10875	-1	92	2	524	1	37	90	3	43	0
Krimson	-1	103	21	520	3	45	81	3	25	0
AGT01	1	76	3	499	1	43	81	3	33	0
CHECK CHARACTERIS		/0	5				02	5		V
Etna	99	1759	58							
	DTM	lbs/ac	site years							

This long-term data is based on results from wide row trials. * Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991. ** () Indicates a variety that is protected by Plant Breeder's Rights legislation that complies with UPOV 1978.

DRY BEANS • YIELDS BY LOCATION • WIDE ROW

		2020 Yield % Check							
Market Class/ Variety	DTM +/- Check	Carman	Morden	Portage	Winkler				
NAVY	+/- T9905		% T	9905					
AAC Shock	-2	79	100	108	95				
Bolt	-2	99	93	96	99				
Indi	-2	97	96	88	120				
AAC Argosy	-1	87	114	122	114				
Armada	0	86	112	72	110				
Nautica	0	79	91	85	87				
T9905	0	100	100	100	100				
HMS Medalist	1	85	97	91	117				
SV1893GH*	1	77	113	107	104				

DRY BEANS + YIELDS BY LOCATION	• WIDE ROW continued				
			2020 Yiel	d % Check	
Market Class/ Variety	DTM +/- Check	Carman	Morden	Portage	Winkler
Varieties that are registered in the	he US or being tested or propose	d for registration in Ca	nada		
15094	-1	99	102	92	116
15095	3	90	96	69	118
CHECK CHARACTERISTICS					
T9905	100	2139	1735	1717	3131
	DTM			s/ac	
	CV %	9.1	11.2	11.7	7.7
	LSD % Sign. Diff.	14 2005	18 yes	18	14
	Seeding Date	yes Jun 6	Jun 3	yes Jun 5	yes Jun 5
	Harvest Date	Sep 29	Sep 14	Oct 8	Oct 15
BLACK	+/- Eclipse	56925	·	clipse	00015
CDC Blackstrap	-5	133	109	133	94
Ace	-2	120	103	125	98
CDC Jet	-1	103	98	92	94
CDC Super Jet	-1	97	76	107	86
Black Tails	0	98	107	110	102
Eclipse**	0	100	100	100	100
CHECK CHARACTERISTICS					
Eclipse	96	1800	1613	1476	3604
	DTM			s/ac	
	CV %	9.1	11.2	11.7	7.7
	LSD %	16	20	21	12
	Sign. Diff.	yes	yes	yes	yes
	Seeding Date	Jun 6	Jun 3	Jun 5	Jun 5
	Harvest Date	Sep 29	Sep 14	Oct 8	Oct 15
PINTO SV6139GR*	+/- of Windbreaker	116		ndbreaker	103
	-2	116	109	103	
Vibrant Windbreaker	-1	122 100	119	102 100	114
Cowboy*	3	100	100	100	100
Monterrey	3	100	105	105	110
La Paz	5	113	116	95	101
	he US or being tested or propose			20	101
16-NP1	4	90	90	80	86
19-382	5	94	94	81	89
ND Palomino*	5	108	100	103	100
19-279	7	100	99	90	97
CHECK CHARACTERISTICS					
Windbreaker	93	2038	2303	2899	3680
	DTM		lbs	s/ac	
	CV %	10.9	6.9	9.3	12.7
	LSD %	20	12	16	-
	Sign. Diff.	yes	yes	yes	no
	Seeding Date	Jun 6	Jun 3	Jun 5	Jun 5
	Harvest Date	Sep 29	Sep 15	Oct 8	Oct 15
GREAT NORTHERN	+/- Pink Panther	120		Panther	110
Aries	-2	130	119	-	118
DARK RED KIDNEY	+/- Pink Panther	100	% Pink 83	Panther _	87
Dynasty Varieties that are registered in th	– he US or being tested or propose			-	0/
161164	5	80	63	_	77
Red Rover	5	54	57	_	55
LIGHT RED KIDNEY	+/- Pink Panther	57		Panther	55
Big Red	+/- FIIK Faittier 0	104	% P IIIK 99	-	85
Pink Panther	0	100	100	_	100
	he US or being tested or propose				
Red Dawn	-9	111	89	_	107
CHECK CHARACTERISTICS					
Pink Panther	99	1699	1924	-	2787
	DTM		lbs	s/ac	
	CV %	13.9	9.6	-	11.9
	LSD %	24	14	-	18
	Sign. Diff.	yes	yes	-	yes
	Seeding Date	Jun 6	Jun 3	Jun 5	Jun 5
	Harvest Date	Sep 29	Sep 21	Oct 8	Oct 15

DRY BEANS • YIELDS BY LOCATION • WIDE ROW continued

		2020 Yield % Check						
Market Class/ Variety	DTM +/- Check	Carman	Morden	Portage	Winkler			
CRANBERRY	+/- Etna	% Etna						
OAC Racer	-1	125	119	-	95			
Etna	0	100	100	-	100			
OAC Candycane	4	128	123	-	146			
Varieties that are registered in th	e US or being tested or propose	ed for registration in Ca	nada					
Amaranto	-3	108	96	-	99			
Krimson	-1	140	132	-	95			
AGT01	1	98	79	-	61			
CHECK CHARACTERISTICS								
Etna	99	1507	1520	-	2401			
	DTM		lbs	/ac				
	CV %	13.9	9.6	-	11.9			
	LSD %	27	18	-	21			
	Sign. Diff.	yes	yes	_	yes			
	Seeding Date	Jun 6	Jun 3	Jun 5	Jun 5			
	Harvest Date	Sep 29	Sep 21	Oct 8	Oct 15			

* 🖗 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

** 🛞 Indicates a variety that is protected by Plant Breeder's Rights legislation that complies with UPOV 1978.

DRY BEANS • YIELDS BY LOCATION • NARROW ROW

					2020 Yield % Check	
Market Class/ Variety	DTM +/- Check	Yield % Check	Site Years Tested	Melita	Morden	Portage
NAVY	+/- CDC Blackstrap	% CDC Blackstrap			% CDC Blackstrap	
HMS Medalist	3	78	3	91	57	86
SV1893GH*	4	88	5	85	75	67
Indi	5	94	7	102	105	88
AAC Argosy	7	90	5	99	93	82
Bolt	7	81	14	102	72	57
AAC Shock	8	84	10	101	81	72
T9905	8	86	12	107	80	67
Nautica	10	85	7	99	57	57
BLACK	+/- CDC Blackstrap	% CDC Blackstrap			% CDC Blackstrap	
CDC Blackstrap*	0	100	23	100	100	100
CDC Jet	5	88	23	110	84	75
CDC Superjet	5	95	23	102	81	80
Eclipse**	6	101	11	114	91	86
CHECK CHARACTERISTICS						
CDC Blackstrap	93	2780	23	3048	3438	3827
	DTM	lbs/ac	site years		lbs/ac	
			CV %	7.7	8.4	9.2
			LSD %	13	12	12
			Sign. Diff.	yes	yes	yes
			Seeding Date	May 19	Jun 3	Jun 5
			Harvest Date	Sep 9	Sep 21	Oct 8
PINTO	+/- Windbreaker	% Windbreaker			% Windbreaker	
SV6139GR*	0	109	8	111	108	130
Windbreaker	0	100	15	100	100	100
Varieties that are registered	in the US or being teste	d or proposed for regist	ration in Canada			
ND Palomino*	1	102	3	111	98	98
CHECK CHARACTERISTICS						
Windbreaker	98	2505	15	3007	3516	2883
	DTM	lbs/ac	site years		lbs/ac	
			CV %	7.7	8.4	9.2
			LSD %	13	12	16
			Sign. Diff.	yes	yes	yes
			Seeding Date	May 19	Jun 3	Jun 5
			Harvest Date	Sep 9	Sep 21	Oct 8

* 🔞 Indicates a variety that is protected by, or has been applied for and pending, Plant Breeder's Rights legislation that complies with UPOV 1991.

** 🛞 Indicates a variety that is protected by Plant Breeder's Rights legislation that complies with UPOV 1978.

Manitoba Pulse and Soybean Buyer List – November 2020

	EDIBLE BEANS	FABA BEANS	LENTILS	PEAS	SOYBEANS			CGC
COMPANY		2	:		:	PHONE	LOCATION	REGULATED
Alliance Pulse Processors Inc. dba AGT Foods Canada	1	<i>✓</i>	\checkmark	1	1	306-525-4490	Regina, SK	<u> </u>
AGT Foods St. Joseph			<i>✓</i>	\checkmark	\checkmark	204-737-2625	St. Joseph, MB	<i>√</i>
All Commodities (AC) Trading Ltd.			\checkmark	\checkmark		204-339-8001	Winnipeg, MB	✓
Avena Foods Ltd. dba Best Booking Pulses Inc			\checkmark	\checkmark		204-857-4451	Portage la Prairie, MB	1
Belle Pulses Ltd.		\checkmark		\checkmark		306-423-5202	Bellevue, SK	~
Besco Grain Ltd.		1		1		204-745-3662	Carman, MB	1
Brett-Young Seeds				~	\checkmark	204-261-7932	Winnipeg, MB	
BroadGrain Commodities Inc.	1	1	1	1	\checkmark	416-504-0070	Toronto, ON	1
C.B. Constantini Ltd.				1		604-669-1212	Vancouver, BC	<i>✓</i>
Cargill Ltd.					1	204-947-6219	Winnipeg, MB	1
Columbia Grain Inc. (CGI) (Walhalla Bean Co.)	1					701-549-3721	Walhalla, ND	1
Delmar Commodities Ltd.	1		1	1	\checkmark	204-331-3696	Winkler, MB	1
ETG Commodities		~	<i>_</i>	1	<i>_</i>	416-900-4148	Mississauga, ON	1
G3 Canada Limited				1		204-983-0239	Winnipeg, MB	1
Gavilon Grain LLC				v	<i>✓</i>			
					V	816-584-2210	Omaha, NB	•
Global Food and Ingredients Inc.		~	1	1		416-840-8590	Toronto, ON	1
Global Grain Canada Ltd.	<i>√</i>					204-829-3641	Plum Coulee, MB	
Hensall District Co-op	<i>√</i>	-	-	\checkmark	-	204-295-3938	Winnipeg, MB	<i></i>
Horizon Agro Inc.	į				 Image: A start of the start of	204-746-2026	Morris, MB	
Kalshea Commodities Inc.	-	-	<i>√</i>	~		204-272-3773	Winnipeg, MB	
Knight Seeds			\checkmark	\checkmark		204-764-2450	Hamiota, MB	
inear Grain Inc.	\checkmark	\checkmark		\checkmark	\checkmark	204-745-6747	Carman, MB	1
ouis Dreyfus Company Canada ULC				\checkmark	✓	403-205-3322	Calgary, AB	1
Marina Commodities Inc.			1	1		204-937-2300	Roblin, MB	1
Masterfeeds		1		~		403-327-2555	Lethbridge, AB	
McDougall Acres Ltd.	1	1	1	1	\checkmark	306-693-3649	Moose Jaw, SK	
Vonsanto					\checkmark	_	Winnipeg, MB	
Natural Proteins Inc.					\checkmark	204-355-5040	Blumenort, MB	
Nu-Vision Commodities				~	1	204-758-3401	St. Jean Baptiste, MB	
Parrheim Foods				1		306-931-1655	Saskatoon, SK	1
Parrish & Heimbecker Ltd.				v _/	1	204-987-4320	Winnipeg, MB	<i>,</i>
Paterson Grain	<i>✓</i>			<i>\</i>	<i></i>	204-956-2090	Winnipeg, MB	V
• FeedMax Corp.	-	-	÷	1		204-523-0682	Killarney, MB	
Pipeline Foods, ULC	÷			1	 Image: A start of the start of	204-594-8750	Winnipeg, MB	√
Prairie Fava Ltd.	-	\checkmark	-		-	204-721-4715	Glenboro, MB	
Providence Grain Group			~	1	\checkmark	780-997-0211	Fort Saskatchewan, AB	<i>√</i>
PS International, LLC DBA Seaboard Special Crops	-	\checkmark	\checkmark	\checkmark		306-565-3934	Regina, SK	1
Richardson International Ltd.				~		204-934-5627	Winnipeg, MB	1
Richardson Pioneer Limited				\checkmark	✓	204-934-5627	Winnipeg, MB	1
Tri Lake Agri Limited				1		204-523-5380	Killarney, MB	1
Roquette Canada Ltd.				1		204-428-3722	Portage la Prairie, MB	1
Rudy Agro Ltd.	1		1	1		306-867-8667	Outlook, SK	1
Scoular Canada Ltd.	1	1	1	1		403-720-9050	Calgary, AB	1
Seed-Ex Inc.				~	 ✓ 	204-737-2000	Letellier, MB	
Semences Prograin Inc.					<i>✓</i>	450-469-5744	Saint-Césaire, QC	
Shafer Commodities Inc.	<i>√</i>	~	1	1	✓ ✓	204-822-6275	Morden, MB	1
	V	v	<i>√</i>	v	v			V
simpson Seeds Inc. Southland Pulse Inc.			✓ ✓	/		306-693-2132	Moose Jaw, SK	
				<i>\</i>		306-634-8008	Estevan, SK	V
The Andersons Inc.			1	<i>√</i>		419-891-6464	Maumee, OH	
/andaele Seeds Ltd.		1		<i>√</i>		204-665-2384	Medora, MB	
/anderveen Commodity Services Ltd.				1	✓	204-745-6444	Carman, MB	1
Viterra Inc.	~		1	1	✓	-	erra sales representative	√
Western Harvest Bean ULC	\checkmark					204-515-7331	Winnipeg, MB	
Wilbur Ellis Company of Canada Ltd.	\checkmark		\checkmark	\checkmark		204-867-8163	Minnedosa, MB	✓
XPT Grain Inc.	1	:	:	\checkmark	:	306-525-0205	Regina, SK	1

The Canada Grain Act requires some elevators and grain dealers to have a Canadian Grain Commission (CGC) license and post security to cover their liabilities (what they owe) to farmers. Grain dealers and operators of primary, terminal and process elevators in western Canada are licensed by the CGC. Seed cleaning plants, which do not purchase grain, and feed mills do not have to be licensed.

It is the responsibility of farmers to satisfy themselves that any company they deal with is financially sound. Questions regarding licencing and security should be directed to the CGC at 800-853-6705 or 204-983-2770.

MPSG's pulse crop buyers list contains the names of companies that have registered with MPSG and are actively purchasing pulse and soybean crops in Manitoba. The word *registered* does not imply endorsement. The complete list is available on our website manitobapulse.ca.