

Soybean Single Inoculant Trial

Trial ID: 2022-S1IN02 – R.M. of MacDonald

Objective: Quantify the agronomic and economic impacts of seed-applied inoculant (single inoculation) vs. no inoculant in soybean fields. This trial requires a minimum field history of three previous soybean crops.

Summary: Nodulation was very similar between treatments. There was no significant yield difference between soybeans with and without a single inoculant. Due to the lack of yield response, there was a decrease in profit/ac in the inoculated area of the trial, equivalent to the cost of the seed-applied inoculant.

Trial	Inform	nation

Treatment	1x LALFIX PROYIELD (liquid on-seed)
Last Soybean Crop	2018
Soybean History	3-yr history
Soil Texture	Clay
Previous Crop	Oats
Tillage	Conventional
Seeding Date	May 26
Variety	DKB008-48
Seeding Rate	183,000 seeds/ac
Row Spacing	20″
Plant Stand @ V1	102,000 plants/ac
Harvest Date	October 19
_	· ·

Precipitation (mm)


	May	Jun	Jul	Aug	Total
Rainfall	144.2	62.8	147.5	86.4	440.9
Normal	58.5	92	77.8	67.6	295.9
% Normal	246%	68%	190%	128%	149%


Nodulation (R2)⁺

	Average nodulation rating	
Single	4.0	
None	3.9	

+ Nodulation is rated on a scale where 0 = no nodules, 1 = poor nodulation (<5/plant), 2 = fair nodulation (<10/plant), 3 = good nodulation (<20/plant) and 4 = excellent nodulation (>20/plant).

NDVI Field Image August 14









Overall Yield & Economics					
	Mean (bu/ac)	Cost ⁺	Change in Profit/ac ⁺⁺		
Single Inoculant	61.1	\$3/ac	-\$3/ac		
Untreated	62.0				
Yield Difference	-0.9				
P-Value	0.3503				
CV	3.3%				
Significance	No	Economic	Νο		

+ Based on an estimated cost of \$3/ac for liquid inoculant; does not include application cost.

++ Yields were not significantly different, therefore profit/ac decreased by the cost/ac of the single inoculation treatment.

